arXiv:math/0510434v3 [math.NT] 8 Jan 2007

REDUCIBILITY OF RATIONAL FUNCTIONS IN SEVERAL VARIABLES

ARNAUD BODIN

Abstract. We prove a analogous of Stein theorem for rational functions in several variables: we bound the number of reducible fibers by a formula depending on the degree of the fraction.

1. Introduction

Let K be an algebraically closed field. Let $f = \frac{p}{q} \in K(\underline{x})$, with $\underline{x} = (x_1, \ldots, x_n)$, $n \geqslant 2$ and $\gcd(p,q) = 1$, the *degree* of f is $\deg f = \max\{\deg p, \deg q\}$. We associate to a fraction $f = \frac{p}{q}$ the pencil $p - \lambda q$, $\lambda \in \hat{K}$ (where we denote $\hat{K} = K \cup \{\infty\}$ and by convention if $\lambda = \infty$ then $p - \lambda q = q$).

For each $\lambda \in \hat{K}$ write the decomposition into irreducible factors:

$$p - \lambda q = \prod_{i=1}^{n_\lambda} F_i^{r_i}.$$

The *spectrum* of f is $\sigma(f) = \{\lambda \in \hat{K} \mid n_\lambda > 1\}$, and the *order of reducibility* is $\rho(f) = \sum_{\lambda \in \hat{K}} (n_\lambda - 1)$.

A fraction f is *composite* if it is the composition of a univariate rational fraction of degree more than 1 with another rational function.

Theorem 1.1. *Let K be an algebraically closed field of characteristic 0. Let $f \in K(\underline{x})$ be non-composite then*

$$\rho(f) < (\deg f)^2 + \deg f.$$

A theorem of Bertini and Krull implies that if f is non-composite then $\sigma(f)$ is finite and we should notice that $\#\sigma(f) \leqslant \rho(f)$. Later on, for an algebraically closed field of characteristic zero and for a polynomial $f \in K[x,y]$, Stein [St] proved the formula $\rho(f) < \deg f$. This formula has been generalized in several directions, see [Na1] for references. For a rational function $f \in \mathbb{C}(x,y)$ a consequence of the work of Ruppert [Ru] on pencil of curves, is that $\#\sigma(f) < (\deg f)^2$. For K algebraically closed (of any characteristic) and $f \in K(x,y)$ Lorenzini

---

Date: February 2, 2008.

[Lo] proved under geometric hypotheses on the pencil $(p - \lambda q)$ that $\rho(f) < (\deg f)^2$. This has been generalized by Vistoli [Vi] for a pencil in several variables for an algebraically closed field of characteristic 0.

Let us give an example extracted from [Lo]. Let $f(x,y) = \frac{x^3+y^3+(1+x+y)^3}{xy(1+x+y)}$, then $\deg(f) = 3$ and $\sigma(f) = \{1, j, j^2, \infty\}$ (where $\{1, j, j^2\}$ are the third roots of unity). For $\lambda \in \sigma(f)$, $(f = \lambda)$ is composed of three lines hence $\rho(f) = 8 = (\deg f)^2 - 1$. Then Lorenzini's bound is optimal in two variables.

The motivation of this work is that we develop the analogous theory of Stein for rational function: composite fractions, kernels of Jacobian derivatives, groups of divisors,... The method for the two variables case is inspired from the work of Stein [St] and the presentation of that work by Najib [Na1]. For completeness even the proofs similar to the ones of Stein have been included. Another motivation is that with a bit more effort we get the case of several variables by following the ideas of [Na1] (see the articles [Na2], [Na3]).

In §2 we prove that a fraction is non-composite if and only its spectrum is finite. Then in §3 we introduce a theory of Jacobian derivation and compute the kernel. Next in §4 we prove that for a non-composite fraction in two variables $\rho(f) < (\deg f)^2 + \deg f$. Finally in §5 we extend this formula to several variables and we end by stating a result for fields of any characteristic.

Acknowledgements: I wish to thank Pierre Dèbes and Salah Najib for discussions and encouragements.

2. Composite rational functions

Let K be an algebraically closed field. Let $\underline{x} = (x_1, \ldots, x_n)$, $n \geqslant 2$.

Definition 2.1. A rational function $f \in K(\underline{x})$ is *composite* if there exist $g \in K(\underline{x})$ and $r \in K(t)$ with $\deg r \geqslant 2$ such that

$$f = r \circ g.$$

Theorem 2.2. *Let $f = \frac{p}{q} \in K(\underline{x})$. The following assertions are equivalent:*

1. *f is composite;*
2. *$p - \lambda q$ is reducible in $K[\underline{x}]$ for all $\lambda \in \hat{K}$ such that* $\deg p - \lambda q = \deg f$*;*
3. *$p - \lambda q$ is reducible in $K[\underline{x}]$ for infinitely many $\lambda \in \hat{K}$.*

Before proving this result we give two corollaries.

Corollary 2.3. *f is non-composite if and only if its spectrum $\sigma(f)$ is finite.*

One aim of this paper is to give a bound for $\sigma(f)$. The hard implication of this theorem (3) $\Rightarrow$ (1) is in fact a reformulation of a theorem of Bertini and Krull.

We also give a nice application pointed out to us by P. Débes:

Corollary 2.4. *Let* $p \in K[\underline{x}]$ *irreducible. Let* $q \in K[\underline{x}]$ *with* $\deg q < \deg p$ *and* $\gcd(p, q) = 1$. *Then for all but finitely many* $\lambda \in K$, $p - \lambda q$ *is irreducible in* $K[\underline{x}]$.

Convention : When we define a fraction $F = \frac{P}{Q}$ we will assume that $\gcd(P, Q) = 1$.

We start with the easy part of Theorem 2.2:

Proof. (2) $\Rightarrow$ (3) is trivial. Let us prove (1) $\Rightarrow$ (2). Let $f = \frac{p}{q}$ be a composite rational function. There exist $g = \frac{u}{v} \in K(\underline{x})$ and $r \in K(t)$ with $k = \deg r \geqslant 2$ such that $f = r \circ g$. Let us write $r = \frac{a}{b}$. Let $\lambda \in \hat{K}$ such that $\deg a - \lambda b = \deg r$ and factorize $a(t) - \lambda b(t) = \alpha(t - t_1)(t - t_2) \cdots (t - t_k)$, $\alpha \in K^*, t_1, \ldots, t_k \in K$. Then

$$p - \lambda q = q \cdot (f - \lambda) = q \cdot \left(\frac{a - \lambda b}{b}\right)(g) = \alpha q \frac{(g - t_1) \cdots (g - t_k)}{b(g)}.$$

Then by multiplication by v^k at the numerator and denominator we get:

$$(p - \lambda q) \cdot (v^k b(g)) = \alpha q(u - t_1 v) \cdots (u - t_k v),$$

which is a polynomial identity. As $\gcd(a, b) = 1$, $\gcd(u, v) = 1$ and $\gcd(p, q) = 1$ then $u - t_1 v, \ldots, u - t_k v$ divide $p - \lambda q$. Hence $p - \lambda q$ is reducible in $K[\underline{x}]$. ☐

Let us reformulate the Bertini-Krull theorem in our context from [Sc, Theorem 37]. It will enable us to end the proof of Theorem 2.2.

Theorem 2.5 (Bertini, Krull)**.** *Let* $F(\underline{x}, \lambda) = p(\underline{x}) - \lambda q(\underline{x}) \in K[\underline{x}, \lambda]$ *an irreducible polynomial. Then the following conditions are equivalent:*

1. $F(\underline{x}, \lambda_0) \in K[\underline{x}]$ *is reducible for all* $\lambda_0 \in K$ *such that* $\deg_{\underline{x}} F(\underline{x}, \lambda_0) = \deg_{\underline{x}} F$.
2. (a) *either there exist* $\phi, \psi \in K[\underline{x}]$ *with* $\deg_{\underline{x}} F > \max\{\deg \phi, \deg \psi\}$, *and* $a_i \in K[\lambda]$, *such that*

$$F(\underline{x}, \lambda) = \sum_{i=0}^{n} a_i(\lambda)\phi(\underline{x})^{n-i}\psi(\underline{x})^i;$$

   (b) *or* $\mathrm{char}(K) = \pi > 0$ *and* $F(\underline{x}, \lambda) \in K[\underline{x}^\pi, \lambda]$, *where* $\underline{x}^\pi = (x_1^\pi, \ldots, x_n^\pi)$.

We now end the proof of Theorem 2.2:

Proof. (3) $\Rightarrow$ (1) Suppose that $p-\lambda_0 q$ is reducible in $K[\underline{x}]$ for infinitely many $\lambda_0 \in \hat{K}$; then it is reducible for all $\lambda_0 \in K$ such that $\deg_{\underline{x}} F(\underline{x},\lambda_0) = \deg_{\underline{x}} F$ (see Corollary 3 of Theorem 32 of [Sc]). We apply Bertini-Krull theorem:

Case (a): $F(\underline{x},\lambda) = p(\underline{x}) - \lambda q(\underline{x})$ can be written:

$$p(\underline{x}) - \lambda q(\underline{x}) = \sum_{i=0}^{n} a_i(\lambda)\phi(\underline{x})^{n-i}\psi(\underline{x})^i.$$

So we may suppose that for $i=1,\ldots,n$, $\deg_\lambda a_i = 1$, let us write $a_i(\lambda) = \alpha_i - \lambda\beta_i$, $\alpha_i,\beta_i \in K$. Then

$$p(\underline{x}) = \sum_{i=0}^{n} \alpha_i \phi(\underline{x})^{n-i}\psi(\underline{x})^i = \phi^n \sum_{i=0}^{n} \alpha_i \Big(\frac{\psi}{\phi}\Big)^i(\underline{x}),$$

and

$$q(\underline{x}) = \sum_{i=0}^{n} \beta_i \phi(\underline{x})^{n-i}\psi(\underline{x})^i = \phi^n \sum_{i=0}^{n} \beta_i \Big(\frac{\psi}{\phi}\Big)^i(\underline{x}).$$

If we set $g(\underline{x}) = \frac{\psi(\underline{x})}{\phi(\underline{x})} \in K[\underline{x}]$, and $r(t) = \frac{\sum_{i=0}^{n}\alpha_i t^i}{\sum_{i=0}^{n}\beta_i t^i}$ then $\frac{p}{q}(\underline{x}) = r\circ g$. Moreover as $\deg_{\underline{x}} F > \max\{\deg\phi, \deg\psi\}$ this implies $n \geqslant 2$ so that $\deg r \geqslant 2$. Then $\frac{p}{q} = f = r\circ g$ is a composite rational function

Case (b): Let $\pi = \operatorname{char}(K) > 0$ and $F(\underline{x},\lambda) = p(\underline{x}) - \lambda q(\underline{x}) \in K[\underline{x}^\pi,\lambda]$, For $\lambda = 0$ it implies that $p(\underline{x}) = P(\underline{x}^\pi)$, then there exists $p' \in K[\underline{x}]$ such that $p(\underline{x}) = (p'(\underline{x}))^\pi$. For $\lambda = -1$ we obtain $s' \in K[\underline{x}]$ such that $p(\underline{x}) + q(\underline{x}) = (s'(\underline{x}))^\pi$. Then $q(\underline{x}) = (p(\underline{x}) + q(\underline{x})) - p(\underline{x}) = (s'(\underline{x}))^\pi - (p'(\underline{x}))^\pi = (s'(\underline{x}) - p'(\underline{x}))^\pi$. Then if we set $q' = s' - p'$ we obtain $q(\underline{x}) = (q'(\underline{x}))^\pi$. Now set $r(t) = t^\pi$ and $g = \frac{p'}{q'}$ we get $f = \frac{p}{q} = \left(\frac{p'}{q'}\right)^\pi = r\circ g$. $\square$

3. Kernel of the Jacobian derivation

We now consider the two variables case and K is an uncountable algebraically closed field of characteristic zero.

3.1. **Jacobian derivation.** Let $f,g \in K(x,y)$, the following formula:

$$D_f(g) = \frac{\partial f}{\partial x}\frac{\partial g}{\partial y} - \frac{\partial f}{\partial y}\frac{\partial g}{\partial x},$$

defines a derivation $D_f : K(x,y) \to K(x,y)$. Notice the $D_f(g)$ is the determinant of the Jacobian matrix of (f,g). We denote by C_f the kernel of D_f:

$$C_f = \{g \in K(x,y) \mid D_f(g) = 0\}.$$

Then C_f is a subfield of $K(x,y)$. We have the inclusion $K(f) \subset C_f$. Moreover if $g^k \in C_f$, $k \in \mathbb{Z} \setminus \{0\}$ then $g \in C_f$.

Lemma 3.1. *Let $f = \frac{p}{q}$, $g \in K(x,y)$. The following conditions are equivalent:*

1. *$g \in C_f$;*
2. *f and g are algebraically dependent;*
3. *g is constant on irreducible components of the curves $(p - \lambda q = 0)$ for all but finitely many $\lambda \in \hat{K}$;*
4. *g is constant on infinitely many irreducible components of the curves $(p - \lambda q = 0)$, $\lambda \in \hat{K}$.*

Corollary 3.2. *If $g \in C_f$ is not a constant then $C_f = C_g$.*

Proof.

- (1) $\Leftrightarrow$ (2). We follow the idea of [Na1] instead of [St]. f and g are algebraically dependent if and only $\mathrm{transc}_K K(f,g) = 1$. And $\mathrm{transc}_K K(f,g) = 1$ if and only the rank of the Jacobian matrix of (f,g) is less or equal to 1, which is equivalent to $g \in C_f$.
- (2) $\Rightarrow$ (3). Let f and g be algebraically dependent. Then there exists a two variables polynomial in f and g that vanishes. Let us write
$$\sum_{i=0}^{n} R_i(f) g^i = 0$$
where $R_i(t) \in K[t]$. Let us write $f = \frac{p}{q}$, $g = \frac{u}{v}$ and $R_n(t) = \alpha(t - \lambda_1)\cdots(t - \lambda_m)$. Then
$$\sum_{i=0}^{n} R_i\Big(\frac{p}{q}\Big)\left(\frac{u}{v}\right)^i = 0, \text{ hence } \sum_{i=0}^{n} R_i\Big(\frac{p}{q}\Big) u^i v^{n-i} = 0.$$
By multiplication by q^d for $d = \max\{\deg R_i\}$ (in order that $q^d R_i(\frac{p}{q})$ are polynomials) we obtain
$$q^d R_n\Big(\frac{p}{q}\Big) u^n = v\left(-q^d R_{n-1}\Big(\frac{p}{q}\Big) u^{n-1} - \cdots\right).$$
As $\gcd(u,v) = 1$ then v divides the polynomial $q^d R_n(\frac{p}{q})$, then v divides $q^{d-m}(p - \lambda_1 q)\cdots(p - \lambda_m q)$. Then all irreducible factors of v divide q or $p - \lambda_i q$, $i = 1, \ldots, m$.

  Let $\lambda \notin \{\infty, \lambda_1, \ldots, \lambda_m\}$. Let V_λ be an irreducible component of $p - \lambda q$, then $V_\lambda \cap Z(v)$ is zero dimensional (or empty). Hence

v is not identically equal to 0 on V_λ. Then for all but finitely many $(x,y) \in V_\lambda$ we get:

$$\sum_{i=0}^{n} R_i(\lambda) g(x,y)^i = 0.$$

Therefore g can only reach a finite number of values $c_1, \ldots, c_n$ (the roots of $\sum_{i=0}^{n} R_i(\lambda)t^i$). Since V_λ is irreducible, g is constant on V_λ.

- (3) $\Rightarrow$ (4). Clear.
- (4) $\Rightarrow$ (1). We first give a proof that if g is constant along an irreducible component V_λ of $(p - \lambda q = 0)$ then $D_f(g) = 0$ on V_λ (we suppose that V_λ is not in the poles of g). Let $(x_0, y_0) \in V_\lambda$ and $t \mapsto p(t)$ be a local parametrization of V_λ around (x_0, y_0). By definition of $p(t)$ we have $f(p(t)) = \lambda$, this implies that:

$$\left\langle \frac{dp}{dt} \mid \overline{\operatorname{grad} f} \right\rangle = \frac{d(f(p(t))}{dt} = 0$$

and by hypotheses g is constant on V_λ this implies $g(p(t))$ is constant and again:

$$\left\langle \frac{dp}{dt} \mid \overline{\operatorname{grad} g} \right\rangle = \frac{d(g(p(t))}{dt} = 0.$$

Then $\operatorname{grad} f$ and $\operatorname{grad} g$ are orthogonal around (x_0, y_0) on V_λ to the same vector, as we are in dimension 2 this implies that the determinant of Jacobian matrix of (f,g) is zero around (x_0, y_0) on V_λ. By extension $D_f(g) = 0$ on V_λ.

We now end the proof: If g is constant on infinitely many irreducible components V_λ of $(p - \lambda q = 0)$ this implies that $D_f(g) = 0$ on infinitely many V_λ. Then $D_f(g) = 0$ in $K(x,y)$.

□

3.2. **Group of the divisors.** Let $f = \frac{p}{q}$, let $\lambda_1, \ldots, \lambda_n \in \hat{K}$, we denote by $G(f; \lambda_1, \ldots, \lambda_n)$ the multiplicative group generated by all the divisors of the polynomials $p - \lambda_i q$, $i = 1, \ldots, n$.

Let

$$d(f) = (\deg f)^2 + \deg f.$$

Lemma 3.3. *Let $F_1, \ldots, F_r \in G(f; \lambda_1, \ldots, \lambda_n)$. If $r \geqslant d(f)$ then there exists a collection of integers $m_1, \ldots, m_r$ (not all equal to zero) such that*

$$g = \prod_{i=1}^{r} F_i^{m_i} \in C_f.$$

Proof. Let $\mu \notin \{\lambda_1, \dots, \lambda_n\}$, and let S be an irreducible component of $(p - \mu q = 0)$. Let $\bar{S}$ be the projective closure of S. The functions F_i restricted to $\bar{S}$ have their poles and zeroes on the points at infinity of S or on the intersection $S \cap Z(F_i) \subset Z(p) \cap Z(q)$.

Let $n : \tilde{S} \to \bar{S}$ be a normalization of $\bar{S}$. The inverse image under normalisation of the points at infinity are denoted by $\{\gamma_1, \dots, \gamma_k\}$, their number verifies $k \leqslant \deg S \leqslant \deg f$.

At a point $\delta \in Z(p) \cap Z(q)$, the number of points of $n^{-1}(\delta)$ is the local number of branches of S at δ then it is less or equal than $\operatorname{ord}_\delta(S)$, where $\operatorname{ord}_\delta(S)$ denotes the order (or multiplicity) of S at δ (see e.g. [Sh], paragraph II.5.3). Then

$$\begin{aligned}\#n^{-1}(\delta) \leqslant \operatorname{ord}_\delta(S) &\leqslant \operatorname{ord}_\delta Z(p-\mu q) \leqslant \operatorname{ord}_\delta Z(p-\mu q) \cdot \operatorname{ord}_\delta Z(p) \\ &\leqslant \operatorname{mult}_\delta(p-\mu q, p) = \operatorname{mult}_\delta(p,q)\end{aligned}$$

where $\operatorname{mult}_\delta(p,q)$ is the intersection multiplicity (see e.g. [Fu]). Then by Bézout theorem:

$$\sum_{\delta \in Z(p)\cap Z(q)} \#n^{-1}(\delta) \leqslant \sum_{\delta \in Z(p)\cap Z(q)} \operatorname{mult}_\delta(p,q) \leqslant \deg p \cdot \deg q \leqslant (\deg f)^2.$$

Then the inverse image under normalisation of $\cup_{i=1}^r S \cap Z(F_i)$ denoted by $\{\gamma_{k+1}, \dots, \gamma_\ell\}$ have less or equal than $(\deg f)^2$ elements. Notice that $\ell \leqslant \deg f + (\deg f)^2 = d(f)$.

Now let ν_{ij} be the order of F_i at γ_j ($i = 1, \dots, r$; $j = 1, \dots, \ell$). Consider the matrix $M = (\nu_{ij})$. Because the degree of the divisor (F_i) (seen over $\tilde{S}$) is zero we get $\sum_{j=1}^{\ell} \nu_{ij} = 0$, for $i = 1, \dots, r$, that means that columns of M are linearly dependent. Then $\operatorname{rk} M < \ell \leqslant d(f)$, by hypothesis $r \geqslant d(f)$, then the rows of M are also linearly dependent. Let $m_1(\mu, S), \dots, m_r(\mu, S)$ such that $\sum_{i=1}^r m_i(\mu, S)\nu_{ij} = 0$, $j = 1, \dots, \ell$.

Consider the function $g_{\mu,S} = \prod_{i=1}^r F_i^{m_i(\lambda, S)}$. Then this function is regular and does not have zeroes or poles at the points γ_j, because $\sum_{i=1}^r m_i(\mu, S)\nu_{ij} = 0$. Then $g_{\mu,S}$ is constant on S.

This construction gives a map $(\mu, S) \mapsto (m_1(\mu, S), \dots, m_r(\mu, S))$ from K to $\mathbb{Z}^r$. Since K is uncountable, there exists infinitely many (μ, S) with the same $(m_1, \dots, m_r)$. Then the function $g = \prod_{i=1}^r F_i^{m_i}$ is constant on infinitely many components of curves of $(p - \mu q = 0)$ and by Lemma 3.1 this implies $g \in C_f$. ☐

3.3. **Non-composite rational function.**

Let $f = \frac{p}{q}$. Let $G(f)$ be the multiplicative group generated by all divisors of the polynomials

$p - \lambda q$ for all $\lambda \in \hat{K}$. In fact we have

$$G(f) = \bigcup_{(\lambda_1,\ldots,\lambda_n)\in K^n} G(f;\lambda_1,\ldots,\lambda_n).$$

Definition 3.4. A family $F_1,\ldots,F_r \in G(f)$ is *f-free* if $(m_1,\ldots,m_r) \in \mathbb{Z}^r$ is such that $\prod_{i=1}^r F_i^{m_i} \in C_f$ then $(m_1,\ldots,m_r) = (0,\ldots,0)$.

A f-free family $F_1,\ldots,F_r \in G(f)$ is *f-maximal* if for all $F \in G(f)$, $\{F_1,\ldots,F_r,F\}$ is not f-free.

Theorem 3.5. *Let $f \in K(x,y)$, $\deg f > 0$. Then the following conditions are equivalent:*

1. $\deg f = \min\{\deg g \mid g \in C_f \setminus K\}$*;*
2. $\sigma(f)$ *is finite;*
3. $C_f = K(f)$*;*
4. f *is non-composite.*

Remark 3.6. This does not give a new proof of "$\sigma(f)$ is finite $\Leftrightarrow$ f is non-composite" because we use Bertini-Krull theorem.

Remark 3.7. The proof (1) $\Rightarrow$ (2) is somewhat easier than in [St], whereas (2) $\Rightarrow$ (3) is more difficult.

Proof.

- (1) $\Rightarrow$ (2). Let us suppose that $\sigma(f)$ is infinite. Set $f = \frac{p}{q}$, with $\gcd(p,q) = 1$. For all $\alpha \in \sigma(f)$, let F_α be an irreducible divisor of $p - \alpha q$, such that $\deg F_\alpha < \deg f$. By Lemma 3.3 there exists a f-maximal family $\{F_1,\ldots,F_r\}$ with $r \leqslant d(f)$. Moreover $r \geqslant 1$ because $\{F_\alpha\}$ is f-free: if not there exists $k \neq 0$ such that $F_\alpha^k \in C_f$ then $F_\alpha \in C_f$, but $\deg F_\alpha < \deg f$ that contradicts the hypothesis of minimality.

  Now the collection $\{F_1,\ldots,F_r,F_\alpha\}$ is not f-free, so that there exist integers $\{m_1(\alpha),\ldots,m_r(\alpha),m(\alpha)\}$, with $m(\alpha) \neq 0$, such that
  $$F_1^{m_1(\alpha)} \cdots F_r^{m_r(\alpha)} \cdot F_\alpha^{m(\alpha)} \in C_f.$$
  Since $\sigma(f)$ is infinite then is equal to $\hat{K}$ minus a finite number of values (see Theorem 2.2) then $\sigma(f)$ is uncountable and the map $\alpha \mapsto (m_1(\alpha),\ldots,m_r(\alpha),m(\alpha))$ is not injective. Let $\alpha \neq \beta$ such that $m_i(\alpha) = m_i(\beta) = m_i$, $i = 1,\ldots,r$ and $m(\alpha) = m(\beta) = m$. Then $F_1^{m_1} \cdots F_r^{m_r} \cdot F_\alpha^m \in C_f$ and $F_1^{m_1} \cdots F_r^{m_r} \cdot F_\beta^m \in C_f$, it implies that $(F_\alpha/F_\beta)^m \in C_f$, therefore $F_\alpha/F_\beta \in C_f$.

  Now $\deg \frac{F_\alpha}{F_\beta} < \deg f$, then by the hypothesis of minimality it proves $\frac{F_\alpha}{F_\beta}$ is a constant. Let $a \in K^*$ such that $F_\alpha = aF_\beta$, by definition F_α divides $p-\alpha q$, but moreover F_α divides $p-\beta q$ (as F_β do). Then as F_α divides both $p-\alpha q$ and $p-\beta q$, F_α divides p and q, that contradicts $\gcd(p,q)=1$.

- (2) $\Rightarrow$ (3). Let $f=\frac{p}{q}$, $\sigma(f)$ finite and $g \in C_f$, we aim at proving that $g \in K(f)$. The proof will be done in several steps:
  (a) *Reduction to the case* $g=\frac{u}{q^\ell}$. Let $g=\frac{u}{v} \in C_f$, then f and g are algebraically dependent, then there exists a polynomial in f and g that vanishes. As before let us write
  $$\sum_{i=0}^{n} R_i(f) g^i = 0$$
  where $R_i(t) \in K[t]$. As $f=\frac{p}{q}$, $g=\frac{u}{v}$ then
  $$\sum_{i=0}^{n} R_i\Big(\frac{p}{q}\Big)\left(\frac{u}{v}\right)^i = 0, \text{ hence } \sum_{i=0}^{n} R_i\Big(\frac{p}{q}\Big)u^i v^{n-i} = 0.$$
  By multiplication by q^d for $d=\max\{\deg R_i\}$ (in order that all $q^d R_i(\frac{p}{q})$ are polynomials) we get:
  $$q^d R_n\Big(\frac{p}{q}\Big)u^n = v\left(-q^d R_{n-1}\Big(\frac{p}{q}\Big)u^{n-1} - \cdots\right).$$
  As $\gcd(u,v)=1$ then v divides the polynomial $q^d R_n(\frac{p}{q})$; we write $vu' = q^d R_n(\frac{p}{q})$ then
  $$g = \frac{u}{v} = \frac{uu'}{q^d R_n(\frac{p}{q})}.$$
  But $R_n(\frac{p}{q}) \in K(\frac{p}{q})$ then $\frac{uu'}{q^d} \in C_f$, but also we have that $g \in K(f)$ if and only if $\frac{uu'}{q^d} \in K(f)$. This proves the reduction.
  (b) *Reduction to the case* $g=qu$. Let $g=\frac{u}{q^\ell} \in C_f$, $\ell \geqslant 0$. As $\sigma(f)$ is finite by Lemma 3.1 we choose $\lambda \in K$ such that $p-\lambda q$ is irreducible and $g \in C_f$ is constant (equal to c) on $p-\lambda q$. As $g=\frac{u}{q^\ell}$,we have $p-\lambda q$ divides $u-cq^\ell$. We can write:
  $$u - cq^\ell = u'(p-\lambda q).$$
  Then
  $$\frac{u}{q^\ell} = \frac{u'}{q^{\ell-1}}\Big(\frac{p}{q}-\lambda\Big) + c.$$

As $\frac{u}{q^\ell}$ and $f = \frac{p}{q}$ are in C_f we get $\frac{u'}{q^{\ell-1}} \in C_f$; moreover $\frac{u}{q^\ell} \in K(f)$ if and only if $\frac{u'}{q^{\ell-1}} \in K(f)$. By induction on $\ell \geqslant 0$ this prove the reduction.

(c) *Reduction to the case $g = q$.* Let $g = qu \in C_f$. g is constant along the irreducible curve $(p - \lambda q = 0)$. Then $qu = u_1(p - \lambda q) + c_1$.

Let $\deg p = \deg q$. Then $q^h u^h = u_1^h(p^h - \lambda q^h)$ (where P^h denotes the homogeneous part of higher degree of the polynomial P). Then $p^h - \lambda q^h$ divides $q^h u^h$ for infinitely many $\lambda \in K$. As $\gcd(p, q) = 1$ this gives a contradiction.

Hence $\deg p \neq \deg q$. We may assume $\deg p > \deg q$ (otherwise $qu \in C_f$ and $\frac{p}{q} \in C_f$ implies $pu \in C_f$). Then we write:

$$qu = qu_1\Big(\frac{p}{q} - \lambda\Big) + c_1,$$

that proves that $qu_1 \in C_f$ and that $qu \in K(f)$ if and only if $qu_1 \in K(f)$. The inequality $\deg p > \deg q$ implies that $\deg u_1 < \deg u$. We continue by induction, $qu_1 = qu_2(\frac{p}{q} - \lambda) + c_2$, with $\deg u_2 < \deg u_1$,..., until we get $\deg u_n = 0$ that is $u_n \in K^*$. Thus we have prove firstly that $qu_n \in C_f$, that is to say $q \in C_f$, and secondly that $qu \in K(f)$ if and only if $q \in K(f)$.

(d) *Case $g = q$.* If $q \in C_f$ then q is constant along the irreducible curve $(p - \lambda q = 0)$ then $q = a(p - \lambda q) + c$, $a \in K^*$. Then

$$q = \frac{c}{1 - a(\frac{p}{q} - \lambda)} \in K\Big(\frac{p}{q}\Big) = K(f).$$

- (3) ⇒ (4). Let us assume that $C_f = K(f)$ and that f is composite, then there exist $r \in K(t)$, $\deg r \geqslant 2$ and $g \in K(x, y)$ such that $f = r \circ g$. By the formula $\deg f = \deg r \cdot \deg g$ we get $\deg f > \deg g$. Now if $r = \frac{a}{b}$ then we have a relation $b(g)f = a(g)$, then f and g are algebraically dependent, hence by Lemma 3.1, $g \in C_f$. As $C_f = K(f)$, there exists $s \in K(t)$ such that $g = s \circ f$. Then $\deg g \geqslant \deg f$. That yields to a contradiction.
- (4) ⇒ (1). Assume that f is non-composite and let $g \in C_f$ of minimal degree. By Corollary 3.2 we get $C_f = C_g$, then $\deg g = \min\{\deg h \mid h \in C_g \setminus K\}$. Then by the already proved implication (1) ⇒ (3) for g, we get $C_g = K(g)$. Then $f \in C_f = C_g = K(g)$, then there exists $r \in K(t)$ such that $f = r \circ g$, but

as f is non-composite then $\deg r = 1$, hence $\deg f = \deg g = \min\{\deg h \mid h \in C_f \setminus K\}$.

□

4. ORDER OF REDUCIBILITY OF RATIONAL FUNCTIONS IN TWO VARIABLES

Let $f = \frac{p}{q} \in K(x,y)$; for all $\lambda \in \hat{K}$, let n_λ be the number of irreducible components of $p - \lambda q$. Let

$$\rho(f) = \sum_{\lambda \in \hat{K}} (n_\lambda - 1).$$

By Theorem 2.2, $\rho(f)$ is finite if and only if f is non-composite. We give a bound for $\rho(f)$. Recall that we defined:

$$d(f) = (\deg f)^2 + \deg f.$$

Theorem 4.1. *Let K be an algebraic closed field of characteristic 0. If $f \in K(x,y)$ is non-composite then*

$$\rho(f) < d(f).$$

Proof. First notice that K can be supposed uncountable, otherwise it can be embedded into an uncountable field L and the spectrum in K would be included in the spectrum in L.

Let us assume that f is non-composite, then by Theorem 2.2 and its corollary we have that $\sigma(f)$ is finite: $\sigma(f) = \{\lambda_1, \dots, \lambda_r\}$. We suppose that $\rho(f) \geqslant d(f)$. Let $f = \frac{p}{q}$. We decompose the polynomials $p - \lambda_i q$ in irreducible factors, for $i = 1, \dots, r$:

$$p - \lambda_i q = \prod_{j=1}^{n_i} F_{i,j}^{k_{i,j}},$$

where n_i stands for n_{λ_i}. Notice that since $\gcd(p,q) = 1$ then $F_{i,j}$ divides $p - \lambda_i q$ but do not divides any of $p - \mu q$, $\mu \neq \lambda_i$. The collection $\{F_{1,1}, \dots, F_{1,n_1-1}, \dots, F_{r,1}, \dots, F_{r,n_r-1}\}$, is included in $G(f, \lambda_1, \dots, \lambda_r)$ and contains $\rho(f) \geqslant d(f)$ elements, then Lemma 3.3 provides a collections $\{m_{1,1}, \dots, m_{1,n_1-1}, \dots, m_{r,1}, \dots, m_{r,n_r-1}\}$ of integers (not all equal to 0) such that

$$g = \prod_{i=1}^{r} \prod_{j=1}^{n_i-1} F_{i,j}^{m_{i,j}} \in C_f. \tag{1}$$

By Theorem 3.5 it implies that $g \in K(f)$, then $g = \frac{u(f)}{v(f)}$, where $u, v \in K[t]$. Let $\mu_1, \dots, \mu_k$ be the roots of u and $\mu_{k+1}, \dots, \mu_\ell$ the roots of v. Then

$$g = \frac{u(\frac{p}{q})}{v(\frac{p}{q})} = \alpha \frac{\prod_{i=1}^{k} \frac{p}{q} - \mu_i}{\prod_{i=k+1}^{\ell} \frac{p}{q} - \mu_i}$$

so that

$$g = \alpha q^{\ell-2k} \frac{\prod_{i=1}^{k} p - \mu_i q}{\prod_{i=k+1}^{\ell} p - \mu_i q}. \tag{2}$$

If $m_{i_0,j_0} \neq 0$ then by the definition of g by equation (1) and by equation (2), we get that F_{i_0,j_0} divides one of the $p - \mu_i q$ or divides q. If F_{i_0,j_0} divides $p - \mu_i q$ then $\mu_i = \lambda_{i_0} \in \sigma(f)$. If F_{i_0,j_0} divides q then $\mu_i = \infty$, so that $\infty \in \sigma(f)$. In both cases $p - \lambda_{i_0} q$ appears in formula (2) at the numerator or at the denominator of g. Then $F_{i_0,n_{i_0}}$ should appears in decomposition (1), that gives a contradiction. Then $\rho(f) < d(f)$. □

5. Extension to several variables

We follows the lines of the proof of [Na3]. We will need a result that claims that the irreducibility and the degree of a family of polynomials remain constant after a generic linear change of coordinates. For $\underline{x} = (x_1, \ldots, x_n)$ and a matrix $B = (b_{ij}) \in Gl_n(K)$, we denote the new coordinates by $B \cdot \underline{x}$:

$$B \cdot \underline{x} = (\sum_{j=1}^{n} b_{1j} x_j, \ldots, \sum_{j=1}^{n} b_{nj} x_j).$$

Proposition 5.1. *Let K be an infinite field. Let $n \geqslant 3$ and $p_1, \ldots, p_\ell \in K[x_1, \ldots, x_n]$ be irreducible polynomials. Then there exists a matrix $B \in Gl_n(K)$ such that for all $i = 1, \ldots, \ell$ we get:*

- *$p_i(B \cdot \underline{x})$ is irreducible in $\overline{K(x_1)}[x_2, \ldots, x_n]$;*
- *$\deg_{(x_2,\ldots,x_n)} p_i(B \cdot \underline{x}) = \deg_{(x_1,\ldots,x_n)} p_i$.*

The proof of this proposition can be derived from [Sm, Ch. 5, Th. 3D] or by using [FJ, Prop. 9.31]. See [Na3] for details.

Now we return to our main result.

Theorem 5.2. *Let K be an algebraically closed field of characteristic 0. Let $f \in K(\underline{x})$ be non-composite then $\rho(f) < (\deg f)^2 + \deg f$.*

Proof. We will prove this theorem by induction on the number n of variables. For $n = 2$, we proved in Theorem 4.1 that $\rho(f) < (\deg f)^2 + \deg f$.

Let $f = \frac{p}{q} \in K(\underline{x})$, with $\underline{x} = (x_1, \dots, x_n)$. We suppose that f is non-composite. For each $\lambda \in \sigma(f)$ we decompose $p - \lambda q$ into irreducible factors:

$$p - \lambda q = \prod_{i=1}^{n_\lambda} F_{\lambda,i}^{r_{\lambda,i}}. \tag{3}$$

We fix $\mu \notin \sigma(f)$. We apply Proposition 5.1 to the polynomials $p - \mu q$ and $F_{\lambda,i}$, for all $\lambda \in \sigma(f)$ and all $i = 1, \dots, n_\lambda$. Then the polynomials $p(B\cdot\underline{x}) - \mu q(B\cdot\underline{x})$ and $F_{\lambda,i}(B\cdot\underline{x})$ are irreducible in $\overline{K(x_1)}[x_2, \dots, x_n]$ and their degrees in $(x_2, \dots, x_n)$ are equals to the degrees in $(x_1, \dots, x_n)$ of $p - \mu q$ and $F_{\lambda,i}$.

Let denote by $k = \overline{K(x_1)}$. This is an uncountable field, algebraically closed of characteristic zero. Now $p(B \cdot \underline{x}) - \mu q(B \cdot \underline{x})$ is irreducible, then $f(B \cdot \underline{x})$ is non-composite in $k(x_2, \dots, x_n)$.

Now equation (3) become:

$$p(B \cdot \underline{x}) - \lambda q(B \cdot \underline{x}) = \prod_{i=1}^{n_\lambda} F_{\lambda,i}(B \cdot \underline{x})^{r_{\lambda,i}}.$$

Which is the decomposition of $p(B \cdot \underline{x}) - \lambda q(B \cdot \underline{x})$ into irreducible factors in $k(x_2, \dots, x_n)$. Then

$$\sigma(f) \subset \sigma(f(B \cdot \underline{x})),$$

where $\sigma(f)$ is a subset of K, and $\sigma(f(B\cdot\underline{x}))$ is a subset of $k = \overline{K(x_1)}$. As n_λ is also the number of distinct irreducible factors of $p(B\cdot\underline{x}) - \lambda q(B\cdot\underline{x})$ we get:

$$\rho(f) \leqslant \rho(f(B \cdot \underline{x})).$$

Now suppose that the result is true for $n - 1$ variables. Then for $f(B \cdot \underline{x}) \in k(x_2, \dots, x_n)$ we get:

$$\rho(f(B \cdot \underline{x})) < (\deg_{(x_2,\dots,x_n)} f(B \cdot \underline{x}))^2 + (\deg_{(x_2,\dots,x_n)} f(B \cdot \underline{x})).$$

Hence:

$$\begin{aligned}
\rho(f) &\leqslant \rho(f(B \cdot \underline{x})) \\
&< (\deg_{(x_2,\dots,x_n)} f(B \cdot \underline{x}))^2 + (\deg_{(x_2,\dots,x_n)} f(B \cdot \underline{x})) \\
&= (\deg_{(x_1,\dots,x_n)} f)^2 + (\deg_{(x_1,\dots,x_n)} f) \\
&= (\deg f)^2 + (\deg f)
\end{aligned}$$

□

If for $n = 2$ we start the induction with Lorenzini's bound $\rho(f) < (\deg f)^2$ we obtain with the same proof the following result for several variables, for K of any characteristic K and a better bound:

Theorem 5.3. *Let K be an algebraically closed field. Let $f \in K(\underline{x})$ be non-composite then $\rho(f) < (\deg f)^2$.*

REFERENCES

[FJ] M. Fried, M. Jarden, *Field arithmetic.* Springer-Verlag, 1986.

[Fu] W. Fulton, *Algebraic curves.* Addison-Wesley, reprint of 1969.

[Lo] D. Lorenzini, *Reducibility of polynomials in two variables.* J. Algebra 156 (1993), 65–75.

[Na1] S. Najib, *Factorisation des polynômes $P(X_1, \ldots, X_n) - \lambda$ et théorème de Stein.* Thesis, University of Lille, 2005.

[Na2] S. Najib, *Sur le spectre d'un polynôme à plusieurs variables.* Acta Arith. 114 (2004), 169–181.

[Na3] S. Najib, *Une généralisation de l'inégalité de Stein-Lorenzini.* J. Algebra 292 (2005), 566-573.

[Ru] W. Ruppert, *Reduzibilität ebener Kurven.* J. Reine Angew. Math. 369 (1986), 167–191.

[Sc] A. Schinzel, *Polynomials with special regard to reducibility.* Encyclopedia of Mathematics and its Applications 77. Cambridge University, 2000.

[Sm] W. Schmidt, *Equations over finite fields. An elementary approach.* Lecture Notes in Mathematics 536. Springer-Verlag, 1976.

[Sh] I. Shafarevich, *Basic algebraic geometry 1.* Springer-Verlag, Berlin, 1994.

[St] Y. Stein, *The total reducibility order of a polynomial in two variables.* Israel J. Math. 68 (1989), 109–122.

[Vi] A. Vistoli, *The number of reducible hypersurfaces in a pencil.* Invent. Math. 112 (1993), 247–262.

LABORATOIRE PAUL PAINLEVÉ, MATHÉMATIQUES, UNIVERSITÉ DE LILLE 1, 59655 VILLENEUVE D'ASCQ, FRANCE.

E-mail address: `Arnaud.Bodin@math.univ-lille1.fr`